arXiv:0706.4008v1 [gr-qc] 27 Jun 2007

Conserved Matter Superenergy Currents for Orthogonally Transitive Abelian G_2 Isometry Groups

Ingemar Eriksson
Matematiska institutionen, Linköpings universitet,
SE-581 83 Linköping, Sweden
ineri@mai.liu.se

June 27, 2007

Abstract

In a previous paper we showed that the electromagnetic superenergy tensor, the Chevreton tensor, gives rise to a conserved current when there is a hypersurface orthogonal Killing vector present. In addition, the current is proportional to the Killing vector. The aim of this paper is to extend this result to the case when we have a two-parameter Abelian isometry group that acts orthogonally transitive on non-null surfaces. It is shown that for four-dimensional Einstein-Maxwell theory with a source-free electromagnetic field, the corresponding superenergy currents lie in the orbits of the group and are conserved. A similar result is also shown to hold for the trace of the Chevreton tensor and for the Bach tensor, and also in Einstein-Klein-Gordon theory for the superenergy of the scalar field. This links up well with the fact that the Bel tensor has these properties and the possibility of constructing conserved mixed currents between the gravitational field and the matter fields.

1 Introduction

In this paper we continue the investigation of conservation laws for the electromagnetic superenergy tensor, the Chevreton tensor, in Einstein-Maxwell spacetimes. In a previous paper [10] we showed that this tensor gives rise to a conserved current whenever there is a hypersurface orthogonal Killing vector present, i.e., if the Killing vector ξ_a satisfies $\xi_{[a}\nabla_b\xi_{c]} = 0$, then

$$H_{abcd}\xi^b\xi^c\xi^d = \omega\xi_a, \qquad \nabla^a(H_{abcd}\xi^b\xi^c\xi^d) = 0, \tag{1}$$

where H_{abcd} is the Chevreton tensor [9, 15],

$$\begin{aligned} H_{abcd} = &-\frac{1}{2}(\nabla_a F_{ce}\nabla_b F_d{}^e + \nabla_b F_{ce}\nabla_a F_d{}^e + \nabla_c F_{ae}\nabla_d F_b{}^e + \nabla_d F_{ae}\nabla_c F_b{}^e) \\ &+\frac{1}{2}(g_{ab}\nabla_f F_{ce}\nabla^f F_d{}^e + g_{cd}\nabla_f F_{ae}\nabla^f F_b{}^e) + \frac{1}{4}(g_{ab}\nabla_c F_{ef}\nabla_d F^{ef} + g_{cd}\nabla_a F_{ef}\nabla_b F^{ef}) \\ &-\frac{1}{4}g_{ab}g_{cd}\nabla_e F_{fg}\nabla^e F^{fg}. \end{aligned} \tag{2}$$

This result holds in four-dimensional Einstein-Maxwell spacetimes with a source-free electromagnetic field, F_{ab}, that inherits the symmetry of the spacetime. A similar situation to when there is a hypersurface orthogonal Killing vector present occurs when there exists a two-parameter isometry group whose surfaces of transitivity are (locally) orthogonal to a family of 2-surfaces. According to Frobenius's theorem the two Killing vectors generating the group, ξ_a and η_a, then satisfies $\xi_{[a}\eta_b\nabla_c\xi_{d]} = 0 = \xi_{[a}\eta_b\nabla_c\eta_{d]}$. We show in this paper, theorem 7, that when the two Killing vectors commute (i.e., the isometry group is Abelian) and form surfaces that are non-null, then the Chevreton tensor again gives rise to conserved currents,

$$H_{abcd}\xi^{Ib}\xi^{Jc}\xi^{Kd} = \omega_{IJK}\xi_a + \Omega_{IJK}\eta_a \qquad \nabla^a(H_{abcd}\xi^{Ib}\xi^{Jc}\xi^{Kd}) = 0, \tag{3}$$

where $I, J, K = 1, 2$ and $\xi^1_a = \xi_a$ and $\xi^2_a = \eta_a$. We also show that similar results hold for the trace of the Chevreton tensor and for the Bach tensor.

This result is interesting not only because it gives conserved quantities for the electromagnetic field, but also because it gives further support to the possibility of creating conserved currents between the electromagnetic field and the gravitational field at the superenergy level.

The Bel-Robinson tensor [2, 3],

$$\begin{aligned} T_{abcd} =& C_{aecf}C_b{}^e{}_d{}^f + C_{aedf}C_b{}^e{}_c{}^f - \frac{1}{2}g_{ab}C_{efcg}C^{ef}{}_d{}^g \\ & - \frac{1}{2}g_{cd}C_{aefg}C_b{}^{efg} + \frac{1}{8}g_{ab}g_{cd}C_{efgh}C^{efgh}, \end{aligned} \tag{4}$$

is a good candidate for representing gravitational energy since it satisfies the Dominant Property [4, 15] and is divergence-free in vacuum. When matter is present, however, neither the Bel-Robinson tensor, nor the Bel tensor,

$$\begin{aligned} B_{abcd} =& R_{aecf}R_b{}^e{}_d{}^f + R_{aedf}R_b{}^e{}_c{}^f - \frac{1}{2}g_{ab}R_{efcg}R^{ef}{}_d{}^g \\ & - \frac{1}{2}g_{cd}R_{aefg}R_b{}^{efg} + \frac{1}{8}g_{ab}g_{cd}R_{efgh}R^{efgh}, \end{aligned} \tag{5}$$

are divergence-free in general. However, there are some cases when it is still possible to construct conserved currents for the gravitational field at the superenergy level. Lazkoz, Senovilla, and Vera [11] have shown that the Bel tensor gives rise to independently conserved currents for general spacetimes when there is a hypersurface orthogonal Killing vector present or when there are two commuting Killing vectors that act orthogonally transitive on non-null surfaces present. In the first case we have the current

$$B_{abcd}\xi^b\xi^c\xi^d = \omega\xi_a, \qquad \nabla^a(B_{abcd}\xi^b\xi^c\xi^d) = 0, \tag{6}$$

and in the second case the four currents

$$B_{a(bcd)}\xi^{Ib}\xi^{Jc}\xi^{Kd} = \omega_{IJK}\xi_a + \Omega_{IJK}\eta_a, \qquad \nabla^a(B_{a(bcd)}\xi^{Ib}\xi^{Jc}\xi^{Kd}) = 0. \tag{7}$$

Also, Senovilla [15] has shown that for Einstein-Klein-Gordon theory, it is possible to construct a mixed conserved superenergy current between the gravitational field and the scalar field when there is a Killing vector present,

$$\nabla^a\left((B_{abcd} + S_{abcd})\xi^b\xi^c\xi^d\right) = 0, \tag{8}$$

where S_{abcd} is the superenergy of the scalar field. When the above isometries are present this breaks up into two separate conserved currents and we show for completeness here and in [10] that the currents constructed from the superenergy of the scalar field also lie in the orbits of the isometry groups.

We hope that it would be possible to construct a similar conserved current between the gravitational field and the electromagnetic field. Senovilla [15] has shown that this is possible in the case of propagation of discontinuities of the fields. For the general case it is not known, but the results of this paper further support that this might be the case. Also, in the spacetime we use as an example of our results, we do have mixed conserved currents.

In the proofs we have opted for expanding the tensors in a basis where the two Killing vectors are taken as two of the basis vectors. It is also possible to take exterior products with the surface element $\xi_{[a}\eta_{b]}$ and using expressions like $2\xi_{[a}\eta_b\nabla_{c]}\xi_d = -\eta_d\xi_{[a}\nabla_b\xi_{c]} + \xi_d\eta_{[a}\nabla_b\xi_{c]}$, but this approach seems to require quite a lot of extra effort.

2 Conventions and some results

We assume that our spacetime is a four-dimensional manifold equipped with a metric of signature -2. We define the Riemann tensor by

$$(\nabla_a\nabla_b - \nabla_b\nabla_a)v_c = -R_{abcd}v^d. \tag{9}$$

The Einstein equations are

$$R_{ab} - \frac{1}{2}Rg_{ab} + \Lambda g_{ab} = -T_{ab}. \tag{10}$$

We will keep the cosmological constant Λ throughout the calculations. If ξ_a is a Killing vector, then $\nabla_a\xi_b = -\nabla_b\xi_a$ and [17],

$$\nabla_a\nabla_b\xi_c = R_{bcad}\xi^d. \tag{11}$$

We also note that the Lie derivative along a Killing vector commutes with the covariant derivative [19],

$$[\pounds_\xi, \nabla_a]T^{b_1\ldots b_i}{}_{c_1\ldots c_j} = 0. \tag{12}$$

We assume that we have two commuting Killing vectors ξ^a and η^a that act orthogonally transitive on non-null surfaces [16],

$$[\xi,\eta] = \xi^b\nabla_b\eta_a - \eta^b\nabla_b\xi_a = 0, \tag{13}$$

$$\xi_{[a}\eta_b\nabla_c\xi_{d]} = \xi_{[a}\eta_b\nabla_c\eta_{d]} = 0, \tag{14}$$

$$\xi_{[a}\eta_{b]}\xi^a\eta^b \neq 0. \tag{15}$$

If we take a basis consisting of ξ_a, η_a, s_a, and t_a, where s_a and t_a are orthogonal to ξ_a and η_a, we can write

$$\nabla_a\xi_b = C_1\xi_{[a}\eta_{b]} + \xi_{[a}(C_2s_{b]} + C_3t_{b]}) + \eta_{[a}(C_4s_{b]} + C_5t_{b]}) + C_6s_{[a}t_{b]}. \tag{16}$$

Taking an exterior product with $\xi_{[c}\eta_{d]}$ gives $C_6 = 0$ and by contracting with $\xi^a\eta^b$ we get

$$\xi^a\eta^b\nabla_a\xi_b = -\xi^a\eta^b\nabla_b\xi_a = -\xi^a\xi^b\nabla_b\eta_a = 0 = C_1\xi_{[a}\eta_{b]}\xi^a\eta^b, \tag{17}$$

so $C_1 = 0$. The structure is the same for $\nabla_a \eta_b$ and we can write

$$\nabla_a \xi_b = 2\xi_{[a} x_{b]} + 2\eta_{[a} y_{b]}, \tag{18}$$

$$\nabla_a \eta_b = 2\xi_{[a} v_{b]} + 2\eta_{[a} w_{b]}, \tag{19}$$

where x_a, y_a, v_a, and w_a are orthogonal to ξ_a and η_a. We will often write this as

$$\nabla_a \xi^I_b = 2\xi_{[a} x^I_{b]} + 2\eta_{[a} y^I_{b]}, \tag{20}$$

where $I = 1, 2$ and $x^1_a = x_a$, $x^2_a = v_a$, $y^1_a = y_a$, and $y^2_a = w_a$. Via Einstein's equations the energy-momentum tensor satisfies [8, 11]

$$\xi_{[a} \eta_b T_{c]d} \xi^d = \xi_{[a} \eta_b T_{c]d} \eta^d = 0, \tag{21}$$

which implies

$$\xi^{Ib} T_{ab} = \alpha^I \xi_a + \beta^I \eta_a, \tag{22}$$

where $\alpha^1 = \alpha$, $\alpha^2 = \gamma$, $\beta^1 = \beta$, and $\beta^2 = \delta$. By taking the Lie derivatives of this equation with respect to ξ_a and η_a we have that

$$\begin{aligned} \xi^a \nabla_a \alpha^I &= 0, & \eta^a \nabla_a \alpha^I &= 0, \\ \xi^a \nabla_a \beta^I &= 0, & \eta^a \nabla_a \beta^I &= 0. \end{aligned} \tag{23}$$

When there is more than one matter field present this, in general, only applies to the total energy-momentum tensor. Here, as well as throughout the text, proportionality factors like α in $\alpha \xi_a$ are generally non-constant scalar functions.

3 Einstein-Klein-Gordon theory

In this section we show that in Einstein-Klein-Gordon theory the superenergy tensor of the scalar (Klein-Gordon) field gives rise conserved currents for Killing vectors that generate an Abelian two-parameter group of isometries that act orthogonally transitive on non-null surfaces. It has previously been shown that the Bel tensor in combination with the superenergy tensor of the scalar field gives rise to conserved currents for Killing vectors [15] and that for this kind of symmetry, or for hypersurface orthogonal Killing vectors, the Bel tensor gives rise to independently conserved currents that lie in the orbits of the group [11]. Hence, the superenergy currents for the scalar field are also independently conserved, and we show here and in [10] for completeness that these currents also lie in the orbits of the group.

The energy-momentum tensor in Einstein-Klein-Gordon theory is given by

$$T_{ab} = -\nabla_a \phi \nabla_b \phi + \frac{1}{2} g_{ab} (\nabla_c \phi \nabla^c \phi + m^2 \phi^2), \tag{24}$$

where the scalar field, ϕ, satisfies the Klein-Gordon equation, $\nabla^c \nabla_c \phi = m^2 \phi$. The superenergy tensor of the scalar field is given by [15]

$$\begin{aligned} S_{abcd} =& \nabla_a \nabla_c \phi \nabla_b \nabla_d \phi + \nabla_a \nabla_d \phi \nabla_b \nabla_c \phi - g_{ab} (\nabla_c \nabla^e \phi \nabla_d \nabla_e \phi + m^2 \nabla_c \phi \nabla_d \phi) \\ & - g_{cd} (\nabla_a \nabla^e \phi \nabla_b \nabla_e \phi + m^2 \nabla_a \phi \nabla_b \phi) \\ & + \frac{1}{2} g_{ab} g_{cd} (\nabla_e \nabla_f \phi \nabla^e \nabla^f \phi + 2m^2 \nabla_e \phi \nabla^e \phi + m^4 \phi^2). \end{aligned} \tag{25}$$

It has the following symmetries, $S_{abcd} = S_{(ab)(cd)} = S_{cdab}$. We can construct the following currents,

$$\begin{aligned}S_{abcd}\xi^{Ib}\xi^{Jc}\xi^{Kd} =& \left(\nabla_a\nabla_c\phi\nabla_b\nabla_d\phi + \nabla_a\nabla_d\phi\nabla_b\nabla_c\phi\right)\xi^{Ib}\xi^{Jc}\xi^{Kd} \\ &- \frac{1}{3}\xi^{Jc}\xi^K_c\left(\nabla_a\nabla^e\phi\nabla_b\nabla_e\phi + m^2\nabla_a\phi\nabla_b\phi\right)\xi^{Ib} + \omega\xi_a + \Omega\eta_a,\end{aligned} \tag{26}$$

where the scalar functions ω and Ω are used to collect proportionality factors of ξ_a and η_a. If the scalar field is massive, $m \neq 0$, it will have a vanishing Lie derivative [15], $\pounds_{\xi^I}\phi = \xi^{Ia}\nabla_a\phi = 0$, so by using Leibniz rule and expanding with (20) we have that

$$\begin{aligned}&\left(\nabla_a\nabla^e\phi\nabla_b\nabla_e\phi + m^2\nabla_a\phi\nabla_b\phi\right)\xi^{Ib} = \nabla_a\nabla^e\phi\nabla_b\nabla_e\phi\xi^{Ib} = \nabla_a\nabla^e\phi\nabla_e\nabla_b\phi\xi^{Ib} \\ =& - \nabla_a\nabla^e\phi\nabla_b\phi\nabla_e\xi^{Ib} = -\xi^e\nabla_a\nabla_e\phi x^{Ib}\nabla_b\phi - \eta^e\nabla_a\nabla_e\phi y^{Ib}\nabla_b\phi \\ =&\nabla_a\xi^e\nabla_e\phi x^{Ib}\nabla_b\phi + \nabla_a\eta^e\nabla_e\phi y^{Ib}\nabla_b\phi = \omega\xi_a + \Omega\eta_a.\end{aligned} \tag{27}$$

For the other type of term present in (26) we similarly have that

$$\nabla_a\nabla_c\phi\nabla_b\nabla_d\phi\xi^{Ib}\xi^{Jc}\xi^{Kd} = -\nabla_a\xi^{Jc}\nabla_c\phi\nabla_b\nabla_d\phi\xi^{Ib}\xi^{Kd} = \omega\xi_a + \Omega\eta_a. \tag{28}$$

Hence, for a massive scalar field the superenergy currents will lie in the orbits of the group, $S_{abcd}\xi^{Ib}\xi^{Jc}\xi^{Kd} = \omega\xi_a + \Omega\eta_a$.

In the massless case, $m = 0$, the scalar field satisfies $\pounds_{\xi^I}\phi = \xi^{Ia}\nabla_a\phi = C_{\xi^I}$, where C_{ξ^I} is a constant. Here we also have from (22) and (24) that $\nabla_a\phi = \alpha'\xi_a + \beta'\eta_a$. The calculations are similar to the massive case and the conclusion the same.

To prove that the currents are divergence-free, we note that the Lie derivative commutes with covariant derivatives for Killing vectors, so the superenergy tensor has vanishing Lie derivative. Since the Killing vectors commute, the Lie derivative of the currents therefore vanishes. We have proven

Theorem 1. *For Einstein-Klein-Gordon spacetimes, possibly with a cosmological constant Λ, which admit an Abelian two-parameter isometry group that act orthogonally transitive on non-null surfaces, the superenergy tensor of the scalar field gives rise to conserved currents that lie in the orbits of the group,*

$$S_{abcd}\xi^{Ib}\xi^{Jc}\xi^{Kd} = \omega_{IJK}\xi_a + \Omega_{IJK}\eta_a, \qquad \nabla^a\left(S_{abcd}\xi^{Ib}\xi^{Jc}\xi^{Kd}\right) = 0, \tag{29}$$

where, in general, the proportionality factors ω_{IJK} and Ω_{IJK} will be non-constant.

Note, we have here assumed a four-dimensional spacetime, but the expansion of (20) is similar in the n-dimensional case and this result thus holds in n dimensions as well.

4 Einstein-Maxwell theory

In this section we will show that if a four-dimensional Einstein-Maxwell spacetime, possibly with a cosmological constant Λ, admits an Abelian two-parameter group of isometries that act orthogonally transitive on non-null surfaces, then the corresponding Chevreton currents constructed from the Killing vectors of the group will lie in the orbits of the group and will be conserved. It is also shown that this also holds for the trace of the Chevreton tensor and for the

Bach tensor. The electromagnetic field is assumed to be source-free. For a null electromagnetic field we will assume that it inherits the symmetries of the spacetime.

The electromagnetic field is described by the Maxwell tensor, $F_{ab} = -F_{ba}$, which in source-free regions satisfies

$$\nabla^a F_{ab} = 0, \qquad \nabla_{[a}F_{bc]} = 0. \tag{30}$$

The energy momentum tensor is given by

$$T_{ab} = -F_{ac}F_b{}^c + \frac{1}{4}g_{ab}F_{cd}F^{cd}. \tag{31}$$

The Ricci scalar, R, satisfies $R = 4\Lambda$, where Λ is the cosmological constant. From (22) we have that

$$\xi^{Ib}F_{ac}F_b{}^c = \alpha'^I \xi_a + \beta'^I \eta_a. \tag{32}$$

Generally, the Lie derivative of the electromagnetic field in four-dimensional Einstein-Maxwell theory satisfies for any Killing vector ξ^a [13, 18]

$$£_\xi F_{ab} = \xi^c \nabla_c F_{ab} + F_{cb}\nabla_a \xi^c + F_{ac}\nabla_b \xi^c = \Psi \overset{*}{F}_{ab}, \tag{33}$$

where $\overset{*}{F}_{ab}$ is the Hodge dual of F_{ab} and Ψ is a constant for non-null fields and satisfies $l_{[a}\nabla_{b]}\Psi = 0$ for null fields, where l_a is the repeated principal null direction of the field.[1] If Ψ is zero, then the electromagnetic field is said to inherit the symmetry of the spacetime. It has been shown that for an Abelian two-parameter group of isometries that acts orthogonally transitive on non-null surfaces, a non-null electromagnetic field inherits those symmetries of the spacetime [13]. In the case of a null electromagnetic field we will assume that it inherits the symmetries. Hence, $£_{\xi^I} F_{ab} = 0$, or

$$\xi^{Ic}\nabla_c F_{ab} = -F_{cb}\nabla_a \xi^{Ic} - F_{ac}\nabla_b \xi^{Ic}. \tag{34}$$

The basic superenergy tensor of the electromagnetic field is given by [15]

$$\begin{aligned} E_{abcd} = &- \nabla_a F_{ce}\nabla_b F_d{}^e - \nabla_b F_{ce}\nabla_a F_d{}^e + g_{ab}\nabla_f F_{ce}\nabla^f F_d{}^e \\ &+ \frac{1}{2}g_{cd}\nabla_a F_{ef}\nabla_b F^{ef} - \frac{1}{4}g_{ab}g_{cd}\nabla_e F_{fg}\nabla^e F^{fg}. \end{aligned} \tag{35}$$

The Chevreton tensor is defined as $H_{abcd} = \frac{1}{2}(E_{abcd} + E_{cdab})$, or

$$\begin{aligned} H_{abcd} = &- \frac{1}{2}(\nabla_a F_{ce}\nabla_b F_d{}^e + \nabla_b F_{ce}\nabla_a F_d{}^e + \nabla_c F_{ae}\nabla_d F_b{}^e + \nabla_d F_{ae}\nabla_c F_b{}^e) \\ &+ \frac{1}{2}(g_{ab}\nabla_f F_{ce}\nabla^f F_d{}^e + g_{cd}\nabla_f F_{ae}\nabla^f F_b{}^e) + \frac{1}{4}(g_{ab}\nabla_c F_{ef}\nabla_d F^{ef} + g_{cd}\nabla_a F_{ef}\nabla_b F^{ef}) \\ &- \frac{1}{4}g_{ab}g_{cd}\nabla_e F_{fg}\nabla^e F^{fg}. \end{aligned} \tag{36}$$

---

[1] It was erroneously stated in [10] that Ψ is always constant, though this did not interfere with the calculations since the electromagnetic field was assumed to inherit the symmetry of the spacetime.

This tensor is completely symmetric in four dimensions, $H_{abcd} = H_{(abcd)}$ [6]. This tensor is more interesting physically than the basic superenergy tensor, because it gives unique currents and a unique divergence and because it shares the symmetries of the Bel tensor. We will now examine the currents that arise when this tensor is contracted with the Killing vectors, ξ^a and η^a, of our two-parameter group. Since the Chevreton tensor is symmetric there are only four different currents, and by interchange of ξ^a and η^a, we only need to consider currents of the form

$$\begin{aligned} H_{abcd}\xi^{Ib}\xi^{Jc}\xi^{Jd} =& -\nabla_a F_{ce}\nabla_b F_d{}^e \xi^{Ib}\xi^{Jc}\xi^{Jd} - \nabla_c F_{ae}\nabla_d F_b{}^e \xi^{Ib}\xi^{Jc}\xi^{Jd} \\ &+ \frac{1}{2}\xi^J_c\xi^{Jc}\nabla_f F_{ae}\nabla^f F_b{}^e\xi^{Ib} + \frac{1}{4}\xi^J_c\xi^{Jc}\nabla_a F_{ef}\nabla_b F^{ef}\xi^{Ib} + \omega\xi_a + \Omega\eta_a, \end{aligned} \tag{37}$$

Here and later ω and Ω are again used to collect the proportionality factors of ξ_a and η_a. We want to show that the remaining terms also lie in the orbits of the group. The proof is divided into three lemmas. We treat the second and third terms separately and then the first and fourth together. The proofs involve some quite lengthy calculations.

Lemma 2. *Under our assumptions,*

$$\xi_{[e}\eta_f\nabla^d F_{a]c}\nabla_d F_b{}^c\xi^{Ib} = 0. \tag{38}$$

Proof. We rewrite $\nabla^d F_{ac}\nabla_d F_b{}^c\xi^{Ib}$ by applying the d'Alembertian $\Box = \nabla^d\nabla_d$ to the energy-momentum tensor (31)

$$\nabla^d F_{ac}\nabla_d F_b{}^c\xi^{Ib} = -\frac{1}{2}\xi^{Ib}(\Box T_{ab} + F_b{}^c\Box F_{ac} + F_a{}^c\Box F_{bc}) + \omega\xi^I_a. \tag{39}$$

By using the four-dimensional Maxwell wave equation, $\Box F_{ab} = 2C^c{}_{ab}{}^d F_{dc} - \frac{1}{3}RF_{ab}$ [14], we get

$$-\frac{1}{2}\xi^{Ib}\Box T_{ab} - \xi^{Ib}(C^d{}_{ac}{}^e F_b{}^c F_{ed} + C^d{}_{bc}{}^e F_a{}^c F_{ed}) + \frac{1}{3}R\xi^{Ib}F_{ac}F_b{}^c + \omega\xi^I_a. \tag{40}$$

By (32), the next to last term equals $\omega\xi_a + \Omega\eta_a$. The two terms involving the Weyl tensor are rewritten using the four-dimensional identity [10, 12]

$$C_{[ab}{}^{[cd}\delta^{f]}_{e]} = 0. \tag{41}$$

We are then left with

$$-\frac{1}{2}\xi^{Ib}\Box T_{ab} - C_{ebad}F^{ec}F_c{}^d\xi^{Ib} + \omega\xi_a + \Omega\eta_a. \tag{42}$$

Substituting the Weyl tensor for Riemann tensor and simplifying with (22) and Einstein's equations (10) yields

$$-\frac{1}{2}\xi^{Ib}\Box T_{ab} - R_{ebad}F^{ec}F_c{}^d\xi^{Ib} + \omega\xi_a + \Omega\eta_a. \tag{43}$$

By (11) we then have that

$$\nabla^d F_{ac}\nabla_d F_b{}^c\xi^{Ib} = -\frac{1}{2}\underbrace{\xi^{Ib}\Box T_{ab}}_{\mathcal{A}} - \underbrace{F^{ec}F_c{}^d\nabla_e\nabla_a\xi^I_d}_{\mathcal{B}} + \omega\xi_a + \Omega\eta_a. \tag{44}$$

We can rewrite the first term, $\mathcal{A}$, with the Leibniz rule as

$$\xi^{Ib}\Box T_{ab} = \Box(T_{ab}\xi^{Ib}) - T_{ab}\Box\xi^{Ib} - 2\nabla_f T_{ab}\nabla^f\xi^{Ib}. \tag{45}$$

For the first term on the right-hand side, use (22), expand, and use (20) and (23). For the second term, use (11) and (22). We are then left with

$$\xi^{Ib}\Box T_{ab} = -2\nabla_f T_{ab}\nabla^f\xi^{Ib} + \omega\xi_a + \Omega\eta_a. \tag{46}$$

Expanding this with (20) and rewriting with the Lie derivative of the energy-momentum tensor, $\pounds_{\xi^I} T_{ab} = \xi^{Ic}\nabla_c T_{ab} + T_{cb}\nabla_a\xi^{Ic} + T_{ac}\nabla_b\xi^{Ic} = 0$, and Leibniz rule, we get

$$2x^{Ib}T_{fb}\nabla_a\xi^f + 2y^{Ib}T_{fb}\nabla_a\eta^f + 2x^{If}\nabla_f(T_{ab}\xi^b) + 2y^{If}\nabla_f(T_{ab}\eta^b) + \omega\xi_a + \Omega\eta_a. \tag{47}$$

Using (22) and expanding again with (20) we are left with only $\omega\xi_a + \Omega\eta_a$.

The second term of (44), $\mathcal{B}$, is expanded with (20) and using (32) we get

$$F^{ec}F_c{}^d\nabla_e\nabla_a\xi^I_d = -(\alpha'\xi^e + \beta'\eta^e)\nabla_e x^I_a - (\gamma'\xi^e + \delta'\eta^e)\nabla_e y^I_a + \omega\xi_a + \Omega\eta_a. \tag{48}$$

Expanding $\pounds_{\xi^I}\nabla_a\xi^J_b = 0$ gives $\pounds_{\xi^I} x^J_b = 0 = \pounds_{\xi^I} y^J_b$, or $\xi^{Ia}\nabla_a x^J_b = \omega\xi_a + \Omega\eta_a$ and $\xi^{Ia}\nabla_a y^J_b = \omega\xi_a + \Omega\eta_a$. Hence,

$$F^{ec}F_c{}^d\nabla_e\nabla_a\xi^I_d = \omega\xi_a + \Omega\eta_a. \tag{49}$$

So, taken together,

$$\xi_{[e}\eta_f\nabla^d F_{a]c}\nabla_d F_b{}^c\xi^{Ib} = 0. \tag{50}$$

□

As in our previous paper [10], we note here that lemma 2 can be applied to the trace of the Chevreton tensor, which is given by [6]

$$H_{ab} = H_{abc}{}^c = \nabla_c F_{ad}\nabla^c F_b{}^d - \frac{1}{4}g_{ab}\nabla_c F_{de}\nabla^c F^{de}. \tag{51}$$

Hence

Theorem 3. *Assume that we have four-dimensional Einstein-Maxwell theory, possibly with a cosmological constant Λ, with a source-free electromagnetic field that inherits the symmetry of the spacetime. If ξ_a and η_a are two commuting Killing vectors that act orthogonally transitive on non-null surfaces, then the currents $H_{ab}\xi^b$ and $H_{ab}\eta^b$, where H_{ab} is the trace of the Chevreton tensor, lie in the orbits of the group,*

$$H_{ab}\xi^b = \omega_1\xi_a + \Omega_1\eta_a, \qquad H_{ab}\eta^b = \omega_2\xi_a + \Omega_2\eta_a, \tag{52}$$

where the proportionality factors ω_i and Ω_i in general are non-constant.

These currents are trivially conserved, since the trace of the Chevreton tensor is divergence-free [6]. Note that for a non-null electromagnetic field we automatically have inherited symmetry.

It was shown in [5] that the trace of the Chevreton tensor is related to the Bach tensor,

$$B_{ab} = \nabla^c \nabla^d C_{acbd} - \frac{1}{2} R^{cd} C_{acbd}. \tag{53}$$

by

$$B_{ab} = 2H_{ab} + \frac{2}{3}\Lambda T_{ab}. \tag{54}$$

Hence, the Bach currents constructed from the Killing vectors ξ_a and η_a will also lie in the orbits of the group. This also applies to the case with a hypersurface orthogonal Killing vector [10].

Corollary 4. *Assume that we have four-dimensional Einstein-Maxwell theory, possibly with a cosmological constant Λ, with a source-free electromagnetic field that inherits the symmetry of the spacetime. If ξ_a is a hypersurface orthogonal Killing vector, then the Bach current $B_{ab}\xi^b$ is proportional to ξ_a,*

$$B_{ab}\xi^b = \omega \xi_a. \tag{55}$$

If ξ_a and η_a are two commuting Killing vectors that act orthogonally transitive on non-null surfaces, then the Bach currents $B_{ab}\xi^b$ and $B_{ab}\eta^b$ lie in the orbits of the group

$$B_{ab}\xi^b = \omega_3 \xi_a + \Omega_3 \eta_a, \qquad B_{ab}\eta^b = \omega_4 \xi_a + \Omega_4 \eta_a. \tag{56}$$

In general, the proportionality factors ω, ω_i, and Ω_i are non-constant.

Again, in the second case, for a non-null electromagnetic field, we automatically have inherited symmetry for those two Killing vectors generating the group.

For the proofs of the following two lemmas we will need to divide into two different cases depending on whether the electromagnetic field is invertible or skew invertible. For a non-null electromagnetic field we can write [13]

$$F_{ab} = \tau_{ab} \cos\alpha + \overset{*}{\tau}_{ab} \sin\alpha, \tag{57}$$

where τ_{ab} is the extremal field and α is the complexion scalar. The extremal field here satisfies one of the following three sets of conditions [13]

$$(1) \qquad \tau_{ab}\xi^a\eta^b = 0 \qquad \overset{*}{\tau}_{ab}\,\xi^a\eta^b = 0, \tag{58}$$

$$(2) \qquad \tau_{ab}\xi^a\eta^b \neq 0 \qquad \overset{*}{\tau}_{ab}\,\xi^a = 0 = \overset{*}{\tau}_{ab}\,\eta^a, \tag{59}$$

$$(3) \qquad \overset{*}{\tau}_{ab}\,\xi^a\eta^b \neq 0 \qquad \tau_{ab}\xi^a = 0 = \tau_{ab}\eta^a. \tag{60}$$

In the first case the electromagnetic field satisfies $F_{ab}\xi^a\eta^b = 0 = \overset{*}{F}_{ab}\,\xi^a\eta^b$, and is said to be skew invertible. It can then be written as

$$F_{ab} = 2\xi_{[a}s_{b]} + 2\eta_{[a}t_{b]}, \tag{61}$$

where s_a and t_a are orthogonal to ξ_a and η_a. Carter [8] showed that the two scalars $F_{ab}\xi^a\eta^b$ and $\overset{*}{F}_{ab}\,\xi^a\eta^b$ are constants and if we, for example, have a spacetime with a symmetry axis where one of the Killing vectors vanishes, the constants vanish everywhere and the electromagnetic field will be skew invertible. In the two other cases the electromagnetic field is invertible and can be written as

$$F_{ab} = 2\kappa\xi_{[a}\eta_{b]} + 2s_{[a}t_{b]}, \tag{62}$$

where again s_a and t_a are orthogonal to ξ_a and η_a.

For a null electromagnetic field with principal null direction l_a we can write

$$F_{ab} = 2l_{[a}A_{b]}, \qquad \overset{*}{F}_{ab}= 2l_{[a}B_{b]}, \tag{63}$$

where A_a and B_a are spacelike vectors satisfying $A^a l_a = B^a l_a = A^a B_a = 0$. By expanding (32) we see that either $l_a = \omega\xi_a + \Omega\eta_a$ or $\xi^a l_a = 0 = \eta^a l_a$, which in either case implies $F_{ab}\xi^a\eta^b = 0 =\overset{*}{F}_{ab}\,\xi^a\eta^b$ and the electromagnetic field is therefore skew invertible.

In the following two proofs we will only show the calculations for the skew invertible case. The invertible case works similarly, noting that $F_{ab}\xi^{Ib} = \lambda^I\xi_a + \mu^I\eta_a$, where $\xi^{Ia}\nabla_a\lambda^J = 0$ and $\xi^{Ia}\nabla_a\mu^J = 0$.

Lemma 5. *Under our assumptions,*

$$\xi_{[f}\eta_g\nabla_{|c|}F_{a]e}\nabla_d F_b{}^e\xi^{Ib}\xi^{Jc}\xi^{Jd} = 0. \tag{64}$$

Proof. We start by rewriting with the Lie derivative (34),

$$\begin{aligned}\nabla_c F_{ae}\nabla_d F_b{}^e\xi^{Ib}\xi^{Jc}\xi^{Jd} =&\xi^{Ib}\nabla_a\xi^{Jc}\left(F_{ce}F_d{}^e\nabla_b\xi^{Jd} + F_{ce}F_{bd}\nabla^e\xi^{Jd}\right)\\ &+ \xi^{Ib}F_{ac}\left(F_d{}^e\nabla_e\xi^{Jc}\nabla_b\xi^{Jd} + F_{bd}\nabla_e\xi^{Jc}\nabla^e\xi^{Jd}\right).\end{aligned} \tag{65}$$

Expanding the first term of the right-hand side with (20) and using (32), we have

$$\xi^{Ib}\nabla_a\xi^{Jc}F_{ce}F_d{}^e\nabla_b\xi^{Jd} = \omega\xi_a + \Omega\eta_a. \tag{66}$$

The second term of (65) is expanded by (20), and if the electromagnetic field is skew invertible, all terms like $F_{ab}\xi^{Ka}\xi^{Lb}$ vanish. Hence,

$$\xi^{Ib}\nabla_a\xi^{Jc}F_{ce}F_{bd}\nabla^e\xi^{Jd} = \omega\xi_a + \Omega\eta_a. \tag{67}$$

For the third term, using (61) and (17) we have

$$\begin{aligned}\xi^{Ib}F_{ac}F_d{}^e\nabla_e\xi^{Jc}\nabla_b\xi^{Jd} &= -s_a\xi_c\xi^{Ib}F_d{}^e\nabla_e\xi^{Jc}\nabla_b\xi^{Jd} - t_a\eta_c\xi^{Ib}F_d{}^e\nabla_e\xi^{Jc}\nabla_b\xi^{Jd} + \omega\xi_a + \Omega\eta_a\\ &= \omega\xi_a + \Omega\eta_a.\end{aligned} \tag{68}$$

For the last term of (65) we expand using (20) and (61) to get

$$\xi^{Ib}F_{ac}F_{bd}\nabla_e\xi^{Jc}\nabla^e\xi^{Jd} = \omega\xi_a + \Omega\eta_a. \tag{69}$$

Hence, taken together, we have that

$$\xi_{[f}\eta_g\nabla_{|c|}F_{a]e}\nabla_d F_b{}^e\xi^{Ib}\xi^{Jc}\xi^{Jd} = 0. \tag{70}$$

The proof is similar for the invertible electromagnetic field.

□

Lemma 6. *Under our assumptions,*

$$-\xi_{[g}\eta_h\nabla_{a]}F_{ce}\nabla_b F_d{}^e\xi^{Ib}\xi^{Jc}\xi^{Jd}+\frac{1}{4}\xi^J_c\xi^{Jc}\xi_{[g}\eta_h\nabla_{a]}F_{ef}\nabla_b F^{ef}\xi^{Ib}=0. \tag{71}$$

Proof. Taking two covariant derivatives of the energy-momentum tensor (31) yields

$$\begin{aligned} -\nabla_a F_{ce}\nabla_b F_d{}^e\xi^{Ib}\xi^{Jc}\xi^{Jd}+\frac{1}{4}\xi^J_c\xi^{Jc}\nabla_a F_{ef}\nabla_b F^{ef}\xi^{Ib} = \\ \frac{1}{2}\underbrace{\nabla_a\nabla_b T_{cd}\xi^{Ib}\xi^{Jc}\xi^{Jd}}_{\mathcal{A}}+\underbrace{F_{ce}\nabla_a\nabla_b F_d{}^e\xi^{Ib}\xi^{Jc}\xi^{Jd}}_{\mathcal{B}}-\frac{1}{4}\xi^{Jc}\xi^J_c\underbrace{\xi^{Ib}F_{ef}\nabla_a\nabla_b F^{ef}}_{\mathcal{C}}. \end{aligned} \tag{72}$$

We rewrite term $\mathcal{A}$ with a covariant derivative of the Lie derivative of the energy-momentum tensor, $\nabla_a \pounds_\xi T_{cd}=0$,

$$\nabla_a\nabla_b T_{cd}\xi^{Ib}\xi^{Jc}\xi^{Jd}=-\nabla_a\xi^{Ib}\nabla_b T_{cd}\xi^{Jc}\xi^{Jd}-2\nabla_a T_{bc}\nabla_d\xi^{Ib}\xi^{Jc}\xi^{Jd}-2T_{bc}\nabla_a\nabla_d\xi^{Ib}\xi^{Jc}\xi^{Jd}. \tag{73}$$

By expanding with (20), the Lie derivative of energy-momentum tensor, (22), and (17),

$$\nabla_a\xi^{Ib}\nabla_b T_{cd}\xi^{Jc}\xi^{Jd}=\omega\xi_a+\Omega\eta_a. \tag{74}$$

Similarly, by using Leibniz rule, expanding with (20) and using (22) and (17)

$$\nabla_a T_{bc}\nabla_d\xi^{Ib}\xi^{Jc}\xi^{Jd}=\omega\xi_a+\Omega\eta_a. \tag{75}$$

Finally, by expanding with (20) twice and using (22), Leibniz rule, and (17),

$$T_{bc}\nabla_a\nabla_d\xi^{Ib}\xi^{Jc}\xi^{Jd}=\omega\xi_a+\Omega\eta_a. \tag{76}$$

Term $\mathcal{B}$, rewritten by taking a covariant derivative of the Lie derivative of the electromagnetic field (34) equals

$$\begin{aligned} F_{ce}\nabla_a\nabla_b F_d{}^e\xi^{Ib}\xi^{Jc}\xi^{Jd}= & -F_c{}^e\nabla_a F_{be}\nabla_d\xi^{Ib}\xi^{Jc}\xi^{Jd}-F_c{}^e\nabla_a F_{db}\nabla_e\xi^{Ib}\xi^{Jc}\xi^{Jd} \\ & -F_c{}^e\nabla_a\xi^{Ib}\nabla_b F_{de}\xi^{Jc}\xi^{Jd}-F_c{}^e F_{be}\nabla_a\nabla_d\xi^{Ib}\xi^{Jc}\xi^{Jd} \end{aligned} \tag{77}$$

The first term is expanded with (20) and if the electromagnetic field is skew invertible, we use (61), to yield

$$-F_c{}^e\nabla_a F_{be}\nabla_d\xi^{Ib}\xi^{Jc}\xi^{Jd}=\omega\xi_a+\Omega\eta_a. \tag{78}$$

The second term is similarly expanded with (20) and (61) and then Leibniz rule is used to give us

$$-F_c{}^e\nabla_a F_{db}\nabla_e\xi^{Ib}\xi^{Jc}\xi^{Jd}=\omega\xi_a+\Omega\eta_a. \tag{79}$$

The third term is expanded with (20) to yield

$$\begin{aligned} -F_c{}^e\nabla_a\xi^{Ib}\nabla_b F_{de}\xi^{Jc}\xi^{Jd} = & x^I_a F_c{}^e\xi^b\nabla_b F_{de}\xi^{Jc}\xi^{Jd}+y^I_a F_c{}^e\eta^b\nabla_b F_{de}\xi^{Jc}\xi^{Jd}+\omega\xi_a+\Omega\eta_a \\ = & \frac{1}{2}x^I_a\xi^b\nabla_b(F_c{}^e F_{de})\xi^{Jc}\xi^{Jd}+\frac{1}{2}y^I_a\eta^b\nabla_b(F_c{}^e F_{de})\xi^{Jc}\xi^{Jd}+\omega\xi_a+\Omega\eta_a, \end{aligned} \tag{80}$$

where, by (32), (23), and (17) we have

$$\begin{aligned}\xi^b\nabla_b(F_c{}^eF_{de})\xi^{Jc}\xi^{Jd} =& \xi^b\nabla_b(F_c{}^eF_{de}\xi^{Jc})\xi^{Jd} - F_c{}^eF_{de}\xi^b\nabla_b\xi^{Jc}\xi^{Jd} \\ =& \xi^b\xi^{Jd}\nabla_b(\alpha'^J\xi_d+\beta'^J\eta_d) - (\alpha'^J\xi_d+\beta'^J\eta_d)\xi^b\nabla_b\xi^{Jd} = 0,\end{aligned} \tag{81}$$

and likewise for the other term. Hence,

$$-F_c{}^e\nabla_a\xi^{Ib}\nabla_bF_{de}\xi^{Jc}\xi^{Jd} = \omega\xi_a+\Omega\eta_a. \tag{82}$$

For the last term we use (32), expand with (20), and Leibniz rule to get

$$-F_c{}^eF_{be}\nabla_a\nabla_d\xi^{Ib}\xi^{Jc}\xi^{Jd} = \omega\xi_a+\Omega\eta_a. \tag{83}$$

Term $\mathcal{C}$, rewritten by taking a covariant derivative of Lie derivative of the electromagnetic field, equals

$$F_{ef}\nabla_a\nabla_bF^{ef}\xi^{Ib} = 2F^{ef}\nabla_aF_{be}\nabla_f\xi^{Ib} - F^{ef}\nabla_a\xi^{Ib}\nabla_bF_{ef} \tag{84}$$

The first term is expanded with (20) and (61) to yield

$$F^{ef}\nabla_aF_{be}\nabla_f\xi^{Ib} = \omega\xi_a+\Omega\eta_a \tag{85}$$

The second term is expanded with (20) and rewritten with the Lie derivative (34),

$$F^{ef}\nabla_a\xi^{Ib}\nabla_bF_{ef} = -x^I_aF^{ef}\xi^b\nabla_bF_{ef} - y^I_aF^{ef}\eta^b\nabla_bF_{ef} + \omega\xi_a+\Omega\eta_a = \omega\xi_a+\Omega\eta_a \tag{86}$$

Hence, taken together, we have that

$$-\xi_{[g}\eta_h\nabla_{a]}F_{ce}\nabla_bF_d{}^e\xi^{Ib}\xi^{Jc}\xi^{Jd} + \frac{1}{4}\xi^J_c\xi^{Jc}\xi_{[g}\eta_h\nabla_{a]}F_{ef}\nabla_bF^{ef}\xi^{Ib} = 0. \tag{87}$$

The case with an invertible electromagnetic field works similarly. □

From (37) together with lemmas 2, 5, and 6 we have that $H_{abcd}\xi^{Ib}\xi^{Jc}\xi^{Kd} = \omega_{IJK}\xi_a + \Omega_{IJK}\eta_a$. The Lie derivative commutes with the covariant derivative for Killing vectors (12), so $\pounds_{\xi^I}\nabla_aF_{bc} = 0$ and we have that $\pounds_{\xi^I}H_{abcd} = 0$. Since the Killing vectors commute, we have that

$$\pounds_{\xi^I}(H_{abcd}\xi^{Jb}\xi^{Kc}\xi^{Ld}) = 0. \tag{88}$$

Hence, the proportionality factors ω and Ω satisfies $\xi^{Ia}\nabla_a\omega = 0 = \xi^{Ia}\nabla_a\Omega$. We have proven

Theorem 7. *Assume that we have four-dimensional Einstein-Maxwell theory, possibly with a cosmological constant, Λ, with a source-free electromagnetic field that inherits the symmetry of the spacetime. If ξ^a and η^a generate a two-parameter Abelian isometry group that act orthogonally transitive on non-null surfaces, then the Chevreton currents constructed from these vectors lie in the orbits of the group and are divergence-free,*

$$\begin{aligned} H_{abcd}\xi^b\xi^c\xi^d &= \omega_5\xi_a+\Omega_5\eta_a, & \nabla^a(H_{abcd}\xi^b\xi^c\xi^d) &= 0.\\ H_{abcd}\xi^b\xi^c\eta^d &= \omega_6\xi_a+\Omega_6\eta_a, & \nabla^a(H_{abcd}\xi^b\xi^c\eta^d) &= 0.\\ H_{abcd}\xi^b\eta^c\eta^d &= \omega_7\xi_a+\Omega_7\eta_a, & \nabla^a(H_{abcd}\xi^b\eta^c\eta^d) &= 0.\\ H_{abcd}\eta^b\eta^c\eta^d &= \omega_8\xi_a+\Omega_8\eta_a, & \nabla^a(H_{abcd}\eta^b\eta^c\eta^d) &= 0. \end{aligned} \tag{89}$$

In general, the proportionality factors ω_i and Ω_i will be non-constant.

Note that for a non-null electromagnetic field, the symmetry is automatically inherited.

In four-dimensional Einstein-Maxwell theory, the Bel tensor (5) can be decomposed as [7, 10]

$$B_{abcd} = T_{abcd} + T_{ab}T_{cd} + \frac{1}{48}R^2 g_{ab}g_{cd}, \tag{90}$$

where T_{abcd} is the Bel-Robinson tensor (4) and T_{ab} is the electromagnetic energy-momentum tensor (31). As was shown in [11], the Bel currents $B_{a(bcd)}\xi^{Ib}\xi^{Jc}\xi^{Kd}$ lie in the orbits of the group. We note that when we contract with the Killing vectors, both the second and the third terms above will lie in the orbits of the group. Both terms also have vanishing Lie derivative, so both terms will give rise to independently conserved currents. From this we see that this also applies to the Bel-Robinson tensor, i.e., it will also give rise to conserved currents that lie in the orbits of the group.

5 Example

Theorem 7 applies to axisymmetric spacetimes, notably the Kerr-Newman solution. The expressions for the currents are however quite large, so we give instead another interesting example – the algebraically general Einstein-Maxwell spacetime found by Barnes [1]. The metric is given by

$$\mathrm{d}s^2 = r\sin(\sqrt{3}\theta)\mathrm{d}x^2 - r\sin(\sqrt{3}\theta)\mathrm{d}y^2 - r2\cos(\sqrt{3}\theta)\mathrm{d}x\mathrm{d}y - r^2\mathrm{d}\theta^2 - \mathrm{d}r^2, \tag{91}$$

and it admits a three-parameter group of isometries generated by the Killing vectors

$$\xi_{1a} = \delta_{xa}, \qquad \xi_{2a} = \delta_{ya}, \qquad \xi_{3a} = y\frac{\sqrt{3}}{2}\delta_{xa} - x\frac{\sqrt{3}}{2}\delta_{ya} + \delta_{\theta a}. \tag{92}$$

Here ξ_1 and ξ_2 form an orthogonally transitive Abelian G_2 subgroup. None of the three Killing vectors are hypersurface orthogonal. The electromagnetic field is given by

$$F_{ab} = 2\cos(p)\delta_{xa}\wedge\delta_{yb} + 2\sin(p)\delta_{ra}\wedge\delta_{\theta b}, \tag{93}$$

where p is an arbitrary constant determining the complexion of the field.

The four Chevreton and Bel currents constructed from ξ_1 and ξ_2 are of course independently conserved. The Chevreton currents are here very simple,

$$\begin{aligned} H_{abcd}\xi_1^b\xi_1^c\xi_1^d &= \frac{3}{2r^2}\xi_{1a}, & H_{abcd}\xi_1^b\xi_1^c\xi_2^d &= \frac{1}{2r^2}\xi_{2a}, \\ H_{abcd}\xi_1^b\xi_2^c\xi_2^d &= \frac{1}{2r^2}\xi_{1a}, & H_{abcd}\xi_2^b\xi_2^c\xi_2^d &= \frac{3}{2r^2}\xi_{2a}. \end{aligned} \tag{94}$$

None of the six Chevreton currents involving the third Killing vector ξ_3 are divergence-free and the same holds for the corresponding Bel currents. However, a combination of the symmetrized Bel tensor with the Chevreton tensor do give conserved currents for all possible combinations of the Killing vectors,

$$\nabla^a\left((B_{a(bcd)} + \frac{1}{3}H_{abcd})\xi_i^b\xi_j^c\xi_k^d\right) = 0. \tag{95}$$

Actually, for this spacetime, all Bel-Robinson currents are independently conserved and it is only the $T_{ab}T_{cd}$ part of the Bel tensor that contributes to the mixed current. For example, the current $v_a = (B_{a(bcd)} + \frac{1}{3}H_{abcd})\xi_1^b\xi_2^c\xi_3^d$, which equals

$$\begin{aligned}
v_x &= -\frac{\sqrt{3}\left(3x\cos(2\sqrt{3}\theta) - 3y\sin(2\sqrt{3}\theta) + 11x\right)}{48r^2},\\
v_y &= \frac{\sqrt{3}\left(3y\cos(2\sqrt{3}\theta) + 3x\sin(2\sqrt{3}\theta) + 11y\right)}{48r^2},\\
v_r &= \frac{\sqrt{3}\sin(\sqrt{3}\theta)}{6r^2},\\
v_\theta &= \frac{\cos(\sqrt{3}\theta)}{3r},
\end{aligned} \tag{96}$$

is conserved. It is also interesting to note that this current is not proportional to a combination of the Killing vectors. This actually holds for all the currents here that involve the third Killing vector ξ_3.

6 Conclusion

We have shown that if a four-dimensional Einstein-Maxwell spacetime admits an Abelian two-parameter isometry group that act orthogonally transitive on non-null surfaces and the electromagnetic field is source-free and inherits the symmetries of the spacetime, then the Chevreton currents generated from the isometry group lies in the orbits of the group and are conserved. Hence, by Gauss's theorem these currents give rise to conserved quantities.

Since the Bel currents have similar properties under this isometry group, this gives further support to the possibility of constructing mixed conserved currents that could govern the interchange of superenergy between the electromagnetic field and the gravitational field.

In the proof of lemma 2 we needed to make use of an identity which holds only in four dimensions, so our result seems to be restricted to this dimension. The results for the Bel currents are n-dimensional, so possible mixed conservation laws may be restricted to four dimensions.

Acknowledgements

The author wishes to thank Göran Bergqvist and José Senovilla for valuable comments and discussions.

References

[1] Barnes A, *A class of algebraically general non-null Einstein-Maxwell fields,* Journal of Physics A: Mathematical and General **9** (1976), 1887–1894.

[2] Bel L, *Sur la radiation gravitionnelle,* Les Comptes Rendus de l'Académie des Sciences de Paris **247** (1958), 1094–1096.

[3] Bel L, *Introduction d'un tenseur de quatriéme ordre,* Les Comptes Rendus de l'Académie des Sciences de Paris **248** (1959), 1297–1300.

[4] Bergqvist G, *Positivity of general superenergy tensors,* Communications in Mathematical Physics **207** (1999), 467–479.

[5] Bergqvist G and Eriksson I, *The Chevreton tensor and Einstein-Maxwell spacetimes conformal to Einstein spaces,* Classical and Quantum Gravity **24** (2007), 3437—3455.

[6] Bergqvist G, Eriksson I, and Senovilla J M M, *New electromagnetic conservation laws,* Classical and Quantum Gravity **20** (2003), 2663–2668.

[7] Bonilla M Á G and Senovilla J M M, *Some properties of the Bel and Bel-Robinson tensors* General Relativity and Gravitation **29** (1997), 91–116.

[8] Carter B, *Killing horizons and orthogonally transitive groups in space-time,* Journal of Mathematical Physics **10** (1969), 70–81.

[9] Chevreton M, *Sur le tenseur de superénergie du champ électromagnétique,* Il Nuovo Cimento **34** (1964), 901–913.

[10] Eriksson I, *Conserved matter superenergy currents for hypersurface orthogonal Killing vectors,* Classical and Quantum Gravity **23** (2006), 2279–2290.

[11] Lazkoz R, Senovilla J M M, and Vera R, *Conserved superenergy currents,* Classical and Quantum Gravity, **20** (2003), 3813–3830.

[12] Lovelock D, *Dimensionally dependent identities,* Mathematical Proceedings of the Cambridge Philosophical Society **68** (1970), 345–350.

[13] Michalski H and Wainwright J, *Killing vector fields and the Einstein-Maxwell field equations in General Relativity,* General Relativity and Gravitation **6** (1975), 289–318.

[14] Penrose R and Rindler W, *Spinors and spacetime vol 2.,* Cambridge University Press, 1986.

[15] Senovilla J M M, *Superenergy tensors,* Classical and Quantum Gravity **17** (2000), 2799–2841.

[16] Stephani H, Kramer D, MacCallum M, Hoenselaers C, and Herlt E, *Exact solutions to Einstein's field equations, 2nd ed.,* Cambridge University Press, 2003.

[17] Wald R M, *General Relativity,* University of Chicago Press, 1984.

[18] Wainwright J and Yaremovicz P A E, *Symmetries and the Einstein-Maxwell field equations - The null field case,* General Relativity and Gravitation **7** (1976), 595–608.

[19] Yano K, *The theory of Lie derivatives and its applications,* North-Holland Publishing Company, 1955.